



[Home](#) / [Investors](#) / [News Releases](#)

# Insmmed to Present New Data from Across its Respiratory Portfolio at the American Thoracic Society 2024 International Conference

*—Additional Data from ARISE Study Evaluating the Impact of ARIKAYCE on Microbiologic Outcomes in Patients with Newly Diagnosed or Recurrent MAC Lung Disease to be Presented as Late-Breaker—*

*—Data to be Presented from ARISE Examining Change in Patient-Reported Respiratory Symptoms—*

BRIDGEWATER, N.J., March 27, 2024 /PRNewswire/ -- Insmmed Incorporated (Nasdaq: INSM), a global biopharmaceutical company on a mission to transform the lives of patients with serious and rare diseases, today announced that it will present nine abstracts from across its respiratory portfolio at the American Thoracic Society (ATS) 2024 International Conference, taking place May 17-22, 2024, in San Diego. Presentations will include late-breaking microbiologic data from the ARISE study of ARIKAYCE® (amikacin liposome inhalation suspension) in patients with newly diagnosed or recurrent *Mycobacterium avium* complex (MAC) lung disease who had not started antibiotics.

"We are excited to present both microbiologic and patient-reported respiratory symptoms data from our Phase 3 ARISE trial of ARIKAYCE in patients with newly diagnosed or recurrent MAC lung disease for the first time in a scientific setting," said Martina Flammer, M.D., M.B.A, Chief Medical Officer of Insmmed. "Further, we look forward to sharing data from across our respiratory portfolio, including brensocatib in bronchiectasis and data on the burden of illness of pulmonary hypertension associated with interstitial lung disease, as we advance our efforts to bring forward novel, urgently needed therapies."

## **Presentations:**

- **Mini Symposium Session A16**, Sunday, May 19, 9:39 AM-9:51 AM PT: [Change in Patient Reported Respiratory Symptoms in a Randomized, Double-blind, Trial of Amikacin Liposome Inhalation Suspension in Adults With Newly Diagnosed or Recurrent \*Mycobacterium Avium\* Complex Lung Disease: The ARISE Study](#)
- **Mini Symposium Session A16 (Late-Breaking Abstract)**, Sunday, May 19, 9:51 AM-10:03 AM PT: [Microbiologic Outcomes From a Randomized, Double-blind Trial of Amikacin Liposome Inhalation Suspension in Adults With Newly Diagnosed or Recurrent \*Mycobacterium Avium\* Complex Lung Disease: The ARISE Study](#)
- **Thematic Poster Session A47**, Sunday, May 19, 11:30 AM-1:15 PM PT:
  - [Incremental Burden of Pulmonary Hypertension Among Patients with Connective Tissue Disease-related Interstitial Lung Disease in the Real-world Setting](#)
  - [Incremental Burden of Pulmonary Hypertension Among Patients with non-Connective Tissue Disease-related Interstitial Lung Disease in the Real-world Setting](#)
- **Thematic Poster Session A74**, Sunday, May 19, 11:30 AM-1:15 PM PT: [Treprostinil Palmitil Conversion Sites in the Lung](#)
- **Poster Discussion Session B26**, Monday, May 20, 9:15 AM-11:15 AM PT: [Incidence and Prevalence of Non-Cystic Fibrosis Bronchiectasis in Japan](#)

- **Poster Discussion Session D25**, Wednesday, May 22, 8:15 AM–10:15 AM PT

- [A Phase 1 Study of Brensocatib Following a Single Oral Administration in Subjects With or Without Renal Impairment](#)
- [Association Between Exacerbation Burden and Comorbidities Among Patients With Non-Cystic Fibrosis Bronchiectasis Over 1 Year](#)

- **Poster Discussion Session D108**, Wednesday, May 22, 11:00 AM—1:00 PM PT: [Novel Anti-inflammatory and Immunomodulatory Effects of the Dipeptidyl Peptidase-1 Inhibitor Brensocatib: A Post-hoc Analysis of the WILLOW Trial](#)

### About ARIKAYCE

ARIKAYCE is approved in the United States as ARIKAYCE<sup>®</sup> (amikacin liposome inhalation suspension), in Europe as ARIKAYCE<sup>®</sup> Liposomal 590 mg Nebuliser Dispersion, and in Japan as ARIKAYCE<sup>®</sup> inhalation 590 mg (amikacin sulfate inhalation drug product). Current international treatment guidelines recommend the use of ARIKAYCE for appropriate patients. ARIKAYCE is a novel, inhaled, once-daily formulation of amikacin, an established antibiotic that was historically administered intravenously and associated with severe toxicity to hearing, balance, and kidney function. Insmed's proprietary PULMOVANCE<sup>®</sup> liposomal technology enables the delivery of amikacin directly to the lungs, where liposomal amikacin is taken up by lung macrophages where the infection resides, while limiting systemic exposure. ARIKAYCE is administered once daily using the Lamira<sup>®</sup> Nebulizer System manufactured by PARI Pharma GmbH (PARI).

### About PARI Pharma and the Lamira<sup>®</sup> Nebulizer System

ARIKAYCE is delivered by a novel inhalation device, the Lamira<sup>®</sup> Nebulizer System, developed by PARI. Lamira<sup>®</sup> is a quiet, portable nebulizer that enables efficient aerosolization of ARIKAYCE via a vibrating, perforated membrane. Based on PARI's 100-year history working with aerosols, PARI is dedicated to advancing inhalation therapies by developing innovative delivery platforms to improve patient care.

### About Brensocatib

Brensocatib is a small molecule, oral, reversible inhibitor of dipeptidyl peptidase 1 (DPP1) being developed by Insmed for the treatment of patients with bronchiectasis, CRSsNP, and other neutrophil-mediated diseases. DPP1 is an enzyme responsible for activating neutrophil serine proteases (NSPs), such as neutrophil elastase, in neutrophils when they are formed in the bone marrow. Neutrophils are the most common type of white blood cell and play an essential role in pathogen destruction and inflammatory mediation. In chronic inflammatory lung diseases, neutrophils accumulate in the airways and result in excessive active NSPs that cause lung destruction and inflammation. Brensocatib may decrease the damaging effects of inflammatory diseases such as bronchiectasis by inhibiting DPP1 and its activation of NSPs. Brensocatib is an investigational drug product that has not been approved for any indication in any jurisdiction.

### About TPIP

Treprostinil palmitil inhalation powder (TPIP) is a dry powder formulation of treprostinil palmitil, a treprostinil prodrug consisting of treprostinil linked by an ester bond to a 16-carbon chain. Developed entirely in Insmed's laboratories, TPIP is a potentially highly differentiated prostanoid being evaluated for the treatment of patients with pulmonary arterial hypertension (PAH), PH-ILD, and other rare and serious pulmonary disorders. TPIP is administered in a capsule-based inhalation device. TPIP is an investigational drug product that has not been approved for any indication in any jurisdiction.

### IMPORTANT SAFETY INFORMATION AND BOXED WARNING FOR ARIKAYCE IN THE U.S.

#### **WARNING: RISK OF INCREASED RESPIRATORY ADVERSE REACTIONS**

**ARIKAYCE has been associated with an increased risk of respiratory adverse reactions, including hypersensitivity pneumonitis, hemoptysis, bronchospasm, and exacerbation of underlying pulmonary disease that have led to hospitalizations in some cases.**

**Hypersensitivity Pneumonitis** has been reported with the use of ARIKAYCE in the clinical trials. Hypersensitivity pneumonitis (reported as allergic alveolitis, pneumonitis, interstitial lung disease, allergic reaction to ARIKAYCE) was reported at a higher frequency in patients treated with ARIKAYCE plus background regimen (3.1%) compared to patients treated with a background regimen alone (0%). Most patients with hypersensitivity pneumonitis discontinued treatment with ARIKAYCE and received treatment with corticosteroids. If hypersensitivity pneumonitis occurs, discontinue ARIKAYCE and manage patients as medically appropriate.

**Hemoptysis** has been reported with the use of ARIKAYCE in the clinical trials. Hemoptysis was reported at a higher frequency in patients treated with ARIKAYCE plus background regimen (17.9%) compared to patients treated with a background regimen alone (12.5%). If hemoptysis occurs, manage patients as medically appropriate.

**Bronchospasm** has been reported with the use of ARIKAYCE in the clinical trials. Bronchospasm (reported as asthma, bronchial hyperreactivity, bronchospasm, dyspnea, dyspnea exertional, prolonged expiration, throat tightness, wheezing) was reported at a higher frequency in patients treated with ARIKAYCE plus background regimen (28.7%) compared to patients treated with a background regimen alone (10.7%). If bronchospasm occurs during the use of ARIKAYCE, treat patients as medically appropriate.

**Exacerbations of underlying pulmonary disease** has been reported with the use of ARIKAYCE in the clinical trials. Exacerbations of underlying pulmonary disease (reported as chronic obstructive pulmonary disease (COPD), infective exacerbation of COPD, infective exacerbation of bronchiectasis) have been reported at a higher frequency in patients treated with ARIKAYCE plus background regimen (14.8%) compared to patients treated with background regimen alone (9.8%). If exacerbations of underlying pulmonary disease occur during the use of ARIKAYCE, treat patients as medically appropriate.

**Anaphylaxis and Hypersensitivity Reactions:** Serious and potentially life-threatening hypersensitivity reactions, including anaphylaxis, have been reported in patients taking ARIKAYCE. Signs and symptoms include acute onset of skin and mucosal tissue hypersensitivity reactions (hives, itching, flushing, swollen lips/tongue/uvula), respiratory difficulty (shortness of breath, wheezing, stridor, cough), gastrointestinal symptoms (nausea, vomiting, diarrhea, crampy abdominal pain), and cardiovascular signs and symptoms of anaphylaxis (tachycardia, low blood pressure, syncope, incontinence, dizziness). Before therapy with ARIKAYCE is instituted, evaluate for previous hypersensitivity reactions to aminoglycosides. If anaphylaxis or a hypersensitivity reaction occurs, discontinue ARIKAYCE and institute appropriate supportive measures.

**Ototoxicity** has been reported with the use of ARIKAYCE in the clinical trials. Ototoxicity (including deafness, dizziness, presyncope, tinnitus, and vertigo) were reported with a higher frequency in patients treated with ARIKAYCE plus background regimen (17%) compared to patients treated with background regimen alone (9.8%). This was primarily driven by tinnitus (7.6% in ARIKAYCE plus background regimen vs 0.9% in the background regimen alone arm) and dizziness (6.3% in ARIKAYCE plus background regimen vs 2.7% in the background regimen alone arm). Closely monitor patients with known or suspected auditory or vestibular dysfunction during treatment with ARIKAYCE. If ototoxicity occurs, manage patients as medically appropriate, including potentially discontinuing ARIKAYCE.

**Nephrotoxicity** was observed during the clinical trials of ARIKAYCE in patients with MAC lung disease but not at a higher frequency than background regimen alone. Nephrotoxicity has been associated with the aminoglycosides. Close monitoring of patients with known or suspected renal dysfunction may be needed when prescribing ARIKAYCE.

**Neuromuscular Blockade:** Patients with neuromuscular disorders were not enrolled in ARIKAYCE clinical trials. Patients with known or suspected neuromuscular disorders, such as myasthenia gravis, should be closely monitored since aminoglycosides may aggravate muscle weakness by blocking the release of acetylcholine at neuromuscular junctions.

**Embryo-Fetal Toxicity:** Aminoglycosides can cause fetal harm when administered to a pregnant woman. Aminoglycosides, including ARIKAYCE, may be associated with total, irreversible, bilateral congenital deafness in pediatric patients exposed *in utero*. Patients who use ARIKAYCE during pregnancy, or become pregnant while taking ARIKAYCE should be apprised of the potential hazard to the fetus.

**Contraindications:** ARIKAYCE is contraindicated in patients with known hypersensitivity to any aminoglycoside.

**Most Common Adverse Reactions:** The most common adverse reactions in Trial 1 at an incidence  $\geq 5\%$  for patients using ARIKAYCE plus background regimen compared to patients treated with background regimen alone were dysphonia (47% vs 1%), cough (39% vs 17%), bronchospasm (29% vs 11%), hemoptysis (18% vs 13%), ototoxicity (17% vs 10%), upper airway irritation (17% vs 2%), musculoskeletal pain (17% vs 8%), fatigue and asthenia (16% vs 10%), exacerbation of underlying pulmonary disease (15% vs 10%), diarrhea (13% vs 5%), nausea (12% vs 4%), pneumonia (10% vs 8%), headache (10% vs 5%), pyrexia (7% vs 5%), vomiting (7% vs 4%), rash (6% vs 2%), decreased weight (6% vs 1%), change in sputum (5% vs 1%), and chest discomfort (5% vs 3%).

**Drug Interactions:** Avoid concomitant use of ARIKAYCE with medications associated with neurotoxicity, nephrotoxicity, and ototoxicity. Some diuretics can enhance aminoglycoside toxicity by altering aminoglycoside concentrations in serum and tissue. Avoid concomitant use of ARIKAYCE with ethacrynic acid, furosemide, urea,

or intravenous mannitol.

**Overdosage:** Adverse reactions specifically associated with overdose of ARIKAYCE have not been identified. Acute toxicity should be treated with immediate withdrawal of ARIKAYCE, and baseline tests of renal function should be undertaken. Hemodialysis may be helpful in removing amikacin from the body. In all cases of suspected overdosage, physicians should contact the Regional Poison Control Center for information about effective treatment.

## U.S. INDICATION

LIMITED POPULATION: ARIKAYCE® is indicated in adults, who have limited or no alternative treatment options, for the treatment of *Mycobacterium avium* complex (MAC) lung disease as part of a combination antibacterial drug regimen in patients who do not achieve negative sputum cultures after a minimum of 6 consecutive months of a multidrug background regimen therapy. As only limited clinical safety and effectiveness data for ARIKAYCE are currently available, reserve ARIKAYCE for use in adults who have limited or no alternative treatment options. This drug is indicated for use in a limited and specific population of patients.

**This indication is approved under accelerated approval based on achieving sputum culture conversion (defined as 3 consecutive negative monthly sputum cultures) by Month 6. Clinical benefit has not yet been established. Continued approval for this indication may be contingent upon verification and description of clinical benefit in confirmatory trials.**

**Limitation of Use:** ARIKAYCE has only been studied in patients with refractory MAC lung disease defined as patients who did not achieve negative sputum cultures after a minimum of 6 consecutive months of a multidrug background regimen therapy. The use of ARIKAYCE is not recommended for patients with non-refractory MAC lung disease.

Patients are encouraged to report negative side effects of prescription drugs to the FDA.

Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch), or call 1-800-FDA-1088. You can also call the Company at 1-844-4-INSMED.

**Please see [Full Prescribing Information](#).**

## About Insmmed

Insmmed Incorporated is a global biopharmaceutical company on a mission to transform the lives of patients with serious and rare diseases. Insmmed's first commercial product is a first-in-disease therapy approved in the United States, Europe, and Japan to treat a chronic, debilitating lung disease. The Company is progressing a robust pipeline of investigational therapies targeting areas of serious unmet need, including neutrophil-mediated inflammatory diseases and rare pulmonary disorders. Insmmed is also advancing an early-stage research engine encompassing a wide range of technologies and modalities, including artificial intelligence-driven protein engineering, gene therapy, and protein manufacturing. Insmmed is headquartered in Bridgewater, New Jersey, with additional offices and research locations throughout the United States, Europe, and Japan. Visit [www.insmed.com](http://www.insmed.com) to learn more.

## Forward-looking Statements

This press release contains forward-looking statements that involve substantial risks and uncertainties.

"Forward-looking statements," as that term is defined in the Private Securities Litigation Reform Act of 1995, are statements that are not historical facts and involve a number of risks and uncertainties. Words herein such as "may," "will," "should," "could," "would," "expects," "plans," "anticipates," "believes," "estimates," "projects," "predicts," "intends," "potential," "continues," and similar expressions (as well as other words or expressions referencing future events, conditions or circumstances) may identify forward-looking statements.

The forward-looking statements in this press release are based upon the Company's current expectations and beliefs, and involve known and unknown risks, uncertainties and other factors, which may cause the Company's actual results, performance and achievements and the timing of certain events to differ materially from the results, performance, achievements or timings discussed, projected, anticipated or indicated in any forward-looking statements. Such risks, uncertainties and other factors include, among others, the following: failure to obtain, or delays in obtaining, regulatory approvals for ARIKAYCE outside the U.S., Europe or Japan, or for the Company's product candidates in the U.S., Europe, Japan or other markets, including separate regulatory approval for the Lamira® Nebulizer System in each market and for each usage; failure to continue to successfully commercialize ARIKAYCE, the Company's only approved product, in the U.S., Europe or Japan (amikacin liposome inhalation suspension, Liposomal 590 mg Nebuliser Dispersion, and amikacin sulfate inhalation drug product, respectively), or to maintain U.S., European or Japanese approval for ARIKAYCE; business or economic disruptions due to catastrophes or other events, including natural disasters or public health crises; risk that brensocatib or TPIP does not prove to be effective or safe for patients in ongoing and future clinical studies, including, for brensocatib, the ASPEN study; uncertainties or changes in the degree of

market acceptance of ARIKAYCE by physicians, patients, third-party payors and others in the healthcare community; the Company's inability to obtain full approval of ARIKAYCE from the FDA, including the risk that the Company will not successfully or in a timely manner validate a PRO tool and complete the confirmatory post-marketing clinical trial required for full approval of ARIKAYCE; inability of the Company, PARI or the Company's other third-party manufacturers to comply with regulatory requirements related to ARIKAYCE or the Lamira<sup>®</sup> Nebulizer System; the Company's inability to obtain adequate reimbursement from government or third-party payors for ARIKAYCE or acceptable prices for ARIKAYCE; development of unexpected safety or efficacy concerns related to ARIKAYCE, brensocatib, TPIP or the Company's other product candidates; inaccuracies in the Company's estimates of the size of the potential markets for ARIKAYCE, brensocatib, TPIP or the Company's other product candidates or in data the Company has used to identify physicians, expected rates of patient uptake, duration of expected treatment, or expected patient adherence or discontinuation rates; the risks and uncertainties associated with, and the perceived benefits of, the Company's secured senior loan with certain funds managed by Pharmakon Advisors, LP and the Company's royalty financing with OrbiMed Royalty & Credit Opportunities IV, LP, including the Company's ability to maintain compliance with the covenants in the agreements for the senior secured loan and royalty financing and the perceived impact of the restrictions on the Company's operations under these agreements; the Company's inability to create or maintain an effective direct sales and marketing infrastructure or to partner with third parties that offer such an infrastructure for distribution of ARIKAYCE or any of the Company's product candidates that are approved in the future; failure to obtain regulatory approval to expand ARIKAYCE's indication to a broader patient population; risk that the Company's competitors may obtain orphan drug exclusivity for a product that is essentially the same as a product the Company is developing for a particular indication; failure to successfully predict the time and cost of development, regulatory approval and commercialization for novel gene therapy products; failure to successfully conduct future clinical trials for ARIKAYCE, brensocatib, TPIP and the Company's other product candidates due to the Company's limited experience in conducting preclinical development activities and clinical trials necessary for regulatory approval and its potential inability to enroll or retain sufficient patients to conduct and complete the trials or generate data necessary for regulatory approval of our product candidates or to permit the use of ARIKAYCE in the broader population of patients with MAC lung disease, among other things; risks that the Company's clinical studies will be delayed, that serious side effects will be identified during drug development, or that any protocol amendments submitted will be rejected; risks that topline, interim or partial data sets are not representative of a complete or larger data set and are subject to change as more patient data becomes available or that blinded data will not be predictive of unblinded data; failure of third parties on which the Company is dependent to manufacture sufficient quantities of ARIKAYCE or the Company's product candidates for commercial or clinical needs, to conduct the Company's clinical trials, or to comply with the Company's agreements or laws and regulations that impact the Company's business or agreements with the Company; the Company's inability to attract and retain key personnel or to effectively manage the Company's growth; the Company's inability to successfully integrate its recent acquisitions and appropriately manage the amount of management's time and attention devoted to integration activities; risks that the Company's acquired technologies, products and product candidates are not commercially successful; the Company's inability to adapt to its highly competitive and changing environment; the Company's inability to access, upgrade or expand its technology systems or difficulties in updating our existing technology or developing or implementing new technology; risk that the Company is unable to maintain its significant customers; risk that government healthcare reform materially increases the Company's costs and damages its financial condition; risk that our current and potential future use of artificial intelligence and machine learning may not be successful; deterioration in general economic conditions in the U.S., Europe, Japan and globally, including the effect of prolonged periods of inflation, affecting the Company, its suppliers, third-party service providers and potential partners; the Company's inability to adequately protect its intellectual property rights or prevent disclosure of its trade secrets and other proprietary information and costs associated with litigation or other proceedings related to such matters; restrictions or other obligations imposed on the Company by agreements related to ARIKAYCE or the Company's product candidates, including its license agreements with PARI and AstraZeneca AB, and failure of the Company to comply with its obligations under such agreements; the cost and potential reputational damage resulting from litigation to which the Company is or may become a party, including product liability claims; risk that the Company's operations are subject to a material disruption in the event of a cybersecurity attack or issue; the Company's limited experience operating internationally; changes in laws and regulations applicable to the Company's business, including any pricing reform, and failure to comply with such laws and regulations; the Company's history of operating losses, and the possibility that the Company may never achieve or maintain profitability; goodwill impairment charges affecting the Company's results of operations and financial condition; inability to repay the Company's existing indebtedness and uncertainties with respect to the Company's ability to access future capital; and delays in the execution of plans to build out an additional third-party manufacturing facility approved by the appropriate regulatory authorities and unexpected expenses associated with those plans.

The Company may not actually achieve the results, plans, intentions or expectations indicated by the Company's forward-looking statements because, by their nature, forward-looking statements involve risks and uncertainties because they relate to events and depend on circumstances that may or may not occur in the

future. For additional information about the risks and uncertainties that may affect the Company's business, please see the factors discussed in Item 1A, "Risk Factors," in the Company's Annual Report on Form 10-K for the year ended December 31, 2023 and any subsequent Company filings with the Securities and Exchange Commission (SEC).

The Company cautions readers not to place undue reliance on any such forward-looking statements, which speak only as of the date of this press release. The Company disclaims any obligation, except as specifically required by law and the rules of the SEC, to publicly update or revise any such statements to reflect any change in expectations or in events, conditions or circumstances on which any such statements may be based, or that may affect the likelihood that actual results will differ from those set forth in the forward-looking statements.

**Contact:**

**Investors:**

Bryan Dunn  
Executive Director, Investor Relations  
Insmed  
(646) 812-4030  
[bryan.dunn@insmed.com](mailto:bryan.dunn@insmed.com)

Eleanor Barisser  
Associate Director, Investor Relations  
Insmed  
(718) 594-5332  
[eleanor.barisser@insmed.com](mailto:eleanor.barisser@insmed.com)

**Media:**

Mandy Fahey  
Executive Director, Corporate Communications  
Insmed  
(732) 718-3621  
[amanda.fahey@insmed.com](mailto:amanda.fahey@insmed.com)

SOURCE Insmed Incorporated

---