



R&D DAY

September 30, 2020

Agenda

Section	Speakers
Welcome & Kick-off	Will Lewis <i>Chair & Chief Executive Officer</i>
Treprostinil Palmitil Inhalation Powder Opportunity	Dr. Eugene Sullivan, M.D. <i>Chief Product Strategy Officer</i> Dr. Ronald J. Oudiz, M.D., FACP, FACC, FCCP <i>Professor of Medicine and Interim Chief Harbor-UCLA Medical Center Division of Cardiology Lead Investigator for Planned Phase 2a Trial</i>
Brensocatib Development Program Update	Dr. Kevin Mange, M.D., MSCE <i>Head of Clinical Development</i> Colleen Silk <i>General Manager, Brensocatib</i>
ARIKAYCE Front Line and Global Expansion Update	Dr. Kevin Mange, M.D., MSCE <i>Head of Clinical Development</i> Neil Hughes <i>General Manager, Head of Europe, Middle East and Africa (EMEA) Commercial</i> Roger Adsett <i>Chief Operating Officer</i>
Financial Update Closing Remarks	Sara Bonstein <i>Chief Financial Officer</i>
Q&A Session	Insmmed Executive Team & Participants



Forward Looking Statements

This presentation contains forward-looking statements that involve substantial risks and uncertainties. "Forward-looking statements," as that term is defined in the Private Securities Litigation Reform Act of 1995, are statements that are not historical facts and involve a number of risks and uncertainties. Words herein such as "may," "will," "should," "could," "would," "expects," "plans," "anticipates," "believes," "estimates," "projects," "predicts," "intends," "potential," "continues," and similar expressions (as well as other words or expressions referencing future events, conditions or circumstances) may identify forward-looking statements.

The forward-looking statements in this presentation are based upon the Company's current expectations and beliefs, and involve known and unknown risks, uncertainties and other factors, which may cause the Company's actual results, performance and achievements and the timing of certain events to differ materially from the results, performance, achievements or timing discussed, projected, anticipated or indicated in any forward-looking statements. Such risks, uncertainties and other factors include, among others, the following: the risk that brensocatib does not prove effective or safe for patients in the STOP-COVID19 study; business or economic disruptions due to catastrophes or other events, including natural disasters or public health crises; impact of the novel coronavirus (COVID-19) pandemic and efforts to reduce its spread on our business, employees, including key personnel, patients, partners and suppliers; failure to successfully commercialize or maintain U.S. approval for ARIKAYCE[®], the Company's only approved product; uncertainties in the degree of market acceptance of ARIKAYCE by physicians, patients, third-party payors and others in the healthcare community; the Company's inability to obtain full approval of ARIKAYCE from the FDA, including the risk that the Company will not timely and successfully complete the study to validate a PRO tool and complete the confirmatory post-marketing study required for full approval of ARIKAYCE; inability of the Company, PARI or the Company's other third party manufacturers to comply with regulatory requirements related to ARIKAYCE or the Lamira[®] Nebulizer System; the Company's inability to obtain adequate reimbursement from government or third-party payors for ARIKAYCE or acceptable prices for ARIKAYCE; development of unexpected safety or efficacy concerns related to ARIKAYCE or brensocatib; inaccuracies in the Company's estimates of the size of the potential markets for ARIKAYCE or its product candidates or in data the Company has used to identify physicians; expected rates of patient uptake, duration of expected treatment, or expected patient adherence or discontinuation rates; the Company's inability to create an effective direct sales

and marketing infrastructure or to partner with third parties that offer such an infrastructure for distribution of ARIKAYCE or any of the Company's product candidates that is approved in the future; failure to obtain regulatory approval to expand ARIKAYCE's indication to a broader patient population; failure to successfully conduct future clinical trials for ARIKAYCE, brensocatib and the Company's other product candidates, including due to the Company's limited experience in conducting preclinical development activities and clinical trials necessary for regulatory approval and the Company's inability to enroll or retain sufficient patients to conduct and complete the trials or generate data necessary for regulatory approval; risks that the Company's clinical studies will be delayed or that serious side effects will be identified during drug development; failure to obtain, or delays in obtaining, regulatory approvals for ARIKAYCE outside the U.S. or for the Company's product candidates in the U.S., Europe, Japan or other markets, including the United Kingdom as a result of its recent exit from the European Union; failure of third parties on which the Company is dependent to manufacture sufficient quantities of ARIKAYCE or the Company's product candidates for commercial or clinical needs, to conduct the Company's clinical trials, or to comply with laws and regulations that impact the Company's business or agreements with the Company; the Company's inability to attract and retain key personnel or to effectively manage the Company's growth; the Company's inability to adapt to its highly competitive and changing environment; the Company's inability to adequately protect its intellectual property rights or prevent disclosure of its trade secrets and other proprietary information and costs associated with litigation or other proceedings related to such matters; restrictions or other obligations imposed on the Company by its agreements related to ARIKAYCE or the Company's product candidates, including its license agreements with PARI and AstraZeneca AB, and failure of the Company to comply with its obligations under such agreements; the cost and potential reputational damage resulting from litigation to which the Company is or may become a party, including product liability claims; the Company's limited experience operating internationally; changes in laws and regulations applicable to the Company's business, including any pricing reform, and failure to comply with such laws and regulations; inability to repay the Company's existing indebtedness and uncertainties with respect to the Company's ability to access future capital; and delays in the execution of plans to build out an additional FDA-approved third-party manufacturing facility and unexpected expenses associated with those plans.

The Company may not actually achieve the results, plans, intentions or

expectations indicated by the Company's forward-looking statements because, by their nature, forward-looking statements involve risks and uncertainties because they relate to events and depend on circumstances that may or may not occur in the future. For additional information about the risks and uncertainties that may affect the Company's business, please see the factors discussed in Item 1A, "Risk Factors," in the Company's Annual Report on Form 10-K for the year ended December 31, 2019, Quarterly Reports on Form 10-Q for the quarters ended March 31, 2020 and June 30, 2020, and any subsequent Company filings with the SEC.

The Company cautions readers not to place undue reliance on any such forward-looking statements, which speak only as of the date of this presentation. The Company disclaims any obligation, except as specifically required by law and the rules of the SEC, to publicly update or revise any such statements to reflect any change in expectations or in events, conditions or circumstances on which any such statements may be based, or that may affect the likelihood that actual results will differ from those set forth in the forward-looking statements.

Will Lewis
Chair & CEO



Mission, Vision & Strategy


Transformational Growth Strategy

World Class Team

Research &
Development Expertise

Global Commercial Presence

Business Development Strength



Leadership Team



William H. Lewis J.D., M.B.A.

Chair & Chief Executive Officer



Roger Adsett, M.B.A.

Chief Operating Officer



Sara M. Bonstein, M.B.A.

Chief Financial Officer



Martina Flammer, M.D., M.B.A.

Chief Medical Officer



Christine Pellizzari, J.D.

Chief Legal Officer



John Soriano, J.D.

Chief Compliance Officer



Nicole Schaffer, M.B.A.

Chief People Strategy Officer



Eugene Sullivan, M.D.

Chief Product Strategy Officer



Neil Hughes, LL.B., M.B.A.

General Manager & Head EMEA



Kevin C. Mange, M.D., MSCE

Head of Clinical Development



Yuji Orihara, M.B.A.

General Manager & Head of Japan



Walter Perkins, Ph.D.

Chief Technology Officer



Colleen Silk, M.B.A.

General Manager - Brensocatib



Andy Stautberg, M.B.A.

General Manager, Early Pipeline



Drayton Wise, M.B.A.

Head of U.S. & General Manager
- ARIKAYCE



Dr. Eugene Sullivan, M.D.

Chief Product Strategy Officer

Dr. Ronald J. Oudiz, M.D., FACP, FACC, FCCP

Professor of Medicine and Interim Chief

Harbor-UCLA Medical Center Division of Cardiology

Lead Investigator for Planned Phase 2a Trial



Treprostinil Palmitil

Development Update

Harnessing the Full Potential of the Prostanoid Pathway in Pulmonary Arterial Hypertension (PAH)

Prostanoids have long been considered the cornerstone therapy for PAH but clinical use has faced challenges related to rapid metabolism and tolerability issues

Treprostinil palmitil has the potential to address the shortcomings of the existing prostanoid therapies by:

- Potentially minimizing side effects, both local and systemic
- Potentially maximizing the vasodilatory activity, by maintaining high local concentrations in the lung
- Allowing for histologic disease-modifying effects of prostanoid therapy

Agenda

1

Pulmonary Arterial Hypertension (PAH)

2

Treprostinil Palmitil Inhalation Powder (TPIP)

3

Highlights of Preclinical Data

4

Product Differentiation

5

Path Forward

Dr. Ronald J. Oudiz, M.D., FACP, FACC, FCCP


Professor of Medicine and Interim Chief

Harbor-UCLA Medical Center Division of Cardiology


Lead Investigator Planned Phase 2a Trial

Pulmonary Arterial Hypertension (PAH) Treatment Landscape


Types of Pulmonary Hypertension




Types of Pulmonary Hypertension



Types of Pulmonary Hypertension



Types of Pulmonary Hypertension



Pulmonary Hypertension: Diagnostic classification

1. Pulmonary arterial hypertension (PAH)

- 1.1. Idiopathic PAH (IPAH)
- 1.2. Heritable
 - 1.2.1. BMPR2
 - 1.2.2. ALK1, ENG, SMAD9, CAV1, KCNK3
 - 1.2.3. Unknown
- 1.3. Drug- and toxin-induced
- 1.4. Associated with
 - 1.4.1. Connective tissue diseases
 - 1.4.2. HIV infection
 - 1.4.3. Portal hypertension
 - 1.4.4. Congenital heart diseases
 - 1.4.5. Schistosomiasis
- 1' Pulmonary veno-occlusive disease and / or pulmonary capillary hemangiomatosis
- 1' Persistent pulmonary hypertension of the newborn

2. Pulmonary hypertension due to left heart disease

- 2.1. Left ventricular systolic dysfunction
- 2.2. Left ventricular diastolic dysfunction
- 2.3. Valvular disease
- 2.4. Congenital/acquired left heart inflow/ outflow tract obstruction and congenital cardiomyopathies

3. PH due to Lung Diseases and/or Hypoxemia

- 3.1. Chronic obstructive pulmonary disease
- 3.2. Interstitial lung disease
- 3.3. Other pulmonary diseases with mixed restrictive and obstructive pattern
- 3.4. Sleep-disordered breathing
- 3.5. Alveolar hypoventilation disorders
- 3.6. Chronic exposure to high altitude
- 3.7. Developmental abnormalities

4. Chronic thromboembolic pulmonary hypertension (CTEPH)

5. Pulmonary hypertension with unclear multifactorial mechanisms

- 5.1. Hematologic disorders: chronic hemolytic anemia, myeloproliferative disorders, splenectomy
- 5.2. Systemic disorders: sarcoidosis, pulmonary histiocytosis: lymphangioleiomyomatosis
- 5.3. Metabolic disorders: glycogen storage disease, Gaucher disease, thyroid disorders
- 5.4. Others: tumoral obstruction, fibrosing mediastinitis, chronic renal failure, segmental PH

Pulmonary Hypertension: Diagnostic classification

1. Pulmonary arterial hypertension (PAH)

- 1.1. Idiopathic PAH (IPAH)
- 1.2. Heritable
 - 1.2.1. BMPR2
 - 1.2.2. ALK1, ENG, SMAD9, CAV1, KCNK3
 - 1.2.3. Unknown
- 1.3. Drug- and toxin-induced
- 1.4. Associated with
 - 1.4.1. Connective tissue diseases
 - 1.4.2. HIV infection
 - 1.4.3. Portal hypertension
 - 1.4.4. Congenital heart diseases
 - 1.4.5. Schistosomiasis
- 1' Pulmonary veno-occlusive disease and / or pulmonary capillary hemangiomatosis
- 1' Persistent pulmonary hypertension of the newborn

2. Pulmonary hypertension due to left heart disease

- 2.1. Left ventricular systolic dysfunction
- 2.2. Left ventricular diastolic dysfunction
- 2.3. Valvular disease
- 2.4. Congenital heart inflow/ outflow tract obstruction
- 2' Right ventricular cardiomyopathies


3. PH due to Lung Diseases and/or Hypoxemia

- 3.1. Chronic obstructive pulmonary disease
- 3.2. Interstitial lung diseases
- 3.3. Other pulmonary diseases with mixed restrictive and obstructive components
- 3.4. Sleep-disordered breathing
- 3.5. Alveolar disorders
- 3.6. Chronic exposure to high altitude
- 3.7. Developmental abnormalities

4. Chronic thromboembolic pulmonary hypertension (CTEPH)


5. Pulmonary hypertension with unclear multifactorial mechanisms

- 5.1. Hematologic disorders: chronic hemolytic anemia, myeloproliferative disorders, splenectomy
- 5.2. Systemic disorders: sarcoidosis, pulmonary histiocytosis: lymphangiomatosis
- 5.3. Metabolic disorders: Gaucher disease, storage disease
- 5.4. Others: fibrosing mediastinitis, congenital aortic valve disease, segmental PH




*D'Alonzo GE, Barst RJ, Ayres SM, et al. Survival in patients with primary pulmonary hypertension. Results from a national prospective registry. *Ann Intern Med* 1991;115:343–349.

Progression of RV Dysfunction




Champion HC, Michelakis ED, Hassoun PM. *Circulation* 2009;120:992-1007


Algorithm for the diagnosis of PH and its causes




Algorithm for the diagnosis of PH and its causes




Algorithm for the diagnosis of PH and its causes (*cont*)



Algorithm for the diagnosis of PH and its causes (*cont*)




Pathways Affecting PAH Treatments




- Endothelin Pathway – targeted by ERAs
- Nitric Oxide Pathway – targeted by PDE5 inhibitors and sGCS
- Prostacyclin Pathway – targeted by prostacyclin analogues and prostacyclin receptor agonists

Pathways Affecting PAH Treatments



- Endothelin Pathway – targeted by ERAs
- Nitric Oxide Pathway – targeted by PDE5 inhibitors and sGCS
- Prostacyclin Pathway – targeted by prostacyclin analogues and prostacyclin receptor agonists

Pathways Affecting PAH Treatments



- Endothelin Pathway – targeted by ERAs
- Nitric Oxide Pathway – targeted by PDE5 inhibitors and sGCS
- Prostacyclin Pathway – targeted by prostacyclin analogues and prostacyclin receptor agonists

Parenteral prostanoids are recommended as first-line therapy for FC IV patients and FC II and III patients with rapid progression. Inhaled prostanoids are recommended for indicated patients unwilling or unable to manage parenteral prostanoids, or as second-line alternatives for patients who are uncontrolled on ERAs or PDE5s (alone or in combination).

PAH Treatments – Approved Therapies

Specific PAH agents

Prostacyclin Analogues

- epoprostenol (IV)
- iloprost (inh)
- treprostinil (IV, SQ, inh, po)
- selexipag (po)

Endothelin antagonists

- bosentan (po)
- ambrisentan (po)
- macitentan (po)

PDE-5 inhibitors

- sildenafil (po)
- tadalafil (po)

Soluble guanylate cyclase agonist

- riociguat (po)

Prostacyclin Advantages and Trade-Offs

Administration	Advantage	Limitations
Oral	Convenient	• GI side effects limit dosing
Parenteral	Effective	• Systemic toxicities limit dosing • Site pain, infection, inconvenience
Inhaled	Targeted	• Short treatment effect requires multiple administrations per day • Sawtooth PK/PD profile

Systemically administered prostanoids (oral or parenteral) are limited by the side effects:

- Hypotension
- Nausea/vomiting
- Diarrhea
- Headache
- Flushing
- Jaw pain

Prostacyclin Advantages and Trade-Offs

Administration	Advantage	Limitations
Oral	Convenient	• GI side effects limit dosing
Parenteral	Effective	• Systemic toxicities limit dosing • Site pain, infection, inconvenience
Inhaled	Targeted	• Short treatment effect requires multiple administrations per day • Sawtooth PK/PD profile

Systemically administered prostanoids (oral or parenteral) are limited by the side effects:

- Hypotension
- Nausea/vomiting
- Diarrhea
- Headache
- Flushing
- Jaw pain

Goal for a new prostanoid therapy: convenient administration; sustained, high lung concentration; lower systemic exposures; potential for anti-proliferative, disease-modifying effects.




Treprostinil Palmitil

**A Potentially
Differentiated Prostanoid**

Treprostinil Palmitil Inhalation Powder (TPIP)

- TPIP is a dry powder formulation of treprostinil prodrug, **treprostinil palmitil***
- The dry powder inhaler (DPI) device is the Plastiape RS01 capsule-based device
- Once delivered to the lung, treprostinil palmitil is hydrolyzed by endogenous esterases to active treprostinil



*Previously treprostinil palmitil was formulated as a nebulized suspension (TPIS).



Highlights of Preclinical Data

Highlights of Preclinical Data

Preclinical data demonstrate the following features of treprostinil palmitil:

Improved pharmacokinetics

- Prolonged lung residence time
- Lower systemic Cmax and prolonged T1/2 compared to treprostinil

Prolonged duration of pharmacodynamic activity compared to treprostinil

“Local” pharmacodynamic effect

- Achieves the same pharmacodynamic (vasodilatory) effect at significantly lower systemic concentrations, compared with IV treprostinil

Effect on hallmark histopathologic features in the sugen/hypoxia model

- Remodeling effect, superior to other prostanoids

Reduced incidence of cough (guinea pig model)

Highlights of Preclinical Data



Potential Implications of Preclinical Data

Preclinical data demonstrate the following features of treprostinil palmitil:

Improved pharmacokinetics

- Prolonged lung residence time
- Lower systemic Cmax and prolonged T1/2 compared to treprostinil

Prolonged duration of pharmacodynamic activity compared to treprostinil

“Local” pharmacodynamic effect

- Achieves the same pharmacodynamic (vasodilatory) effect at significantly lower systemic concentrations, compared with IV treprostinil

Effect on hallmark histopathologic features in the sugen/hypoxia model

- Remodeling effect, superior to other prostanoids

Reduced incidence of cough (guinea pig model)

Potential Implications of Preclinical Data

Preclinical data demonstrate the following features of treprostinil palmitil:

Once-or-twice-daily administration

Improved pharmacokinetics

- Prolonged lung residence time
- Lower systemic C_{max} and prolonged T_{1/2} compared to treprostinil

Prolonged duration of pharmacodynamic activity compared to treprostinil

“Local” pharmacodynamic effect

- Achieves the same pharmacodynamic (vasodilatory) effect at significantly lower systemic concentrations, compared with IV treprostinil

Effect on hallmark histopathologic features in the sugen/hypoxia model

- Remodeling effect, superior to other prostanoids

Reduced incidence of cough (guinea pig model)

Potential Implications of Preclinical Data

Preclinical data demonstrate the following features of treprostinil palmitil:

Comparable effects at lower systemic exposures

Improved pharmacokinetics

- **Prolonged lung residence time**
- Lower systemic C_{max} and prolonged T_{1/2} compared to treprostinil

Prolonged duration of pharmacodynamic activity compared to treprostinil

“Local” pharmacodynamic effect

- Achieves the same pharmacodynamic (vasodilatory) effect at significantly lower systemic concentrations, compared with IV treprostinil

Effect on hallmark histopathologic features in the sugen/hypoxia model

- Remodeling effect, superior to other prostanoids

Reduced incidence of cough (guinea pig model)

Potential Implications of Preclinical Data

Preclinical data demonstrate the following features of treprostinil palmitil:

Potential disease modifying activity

Improved pharmacokinetics

- Prolonged lung residence time
- Lower systemic Cmax and prolonged T1/2 compared to treprostinil

Prolonged duration of pharmacodynamic activity compared to treprostinil

“Local” pharmacodynamic effect

- Achieves the same pharmacodynamic (vasodilatory) effect at significantly lower systemic concentrations, compared with IV treprostinil

Effect on hallmark histopathologic features in the sugen/hypoxia model

- Remodeling effect, superior to other prostanoids

Reduced incidence of cough (guinea pig model)

Potential Implications of Preclinical Data

Preclinical data demonstrate the following features of treprostinil palmitil:

Potential Improved Tolerability

Improved pharmacokinetics

- Prolonged lung residence time
- **Lower systemic C_{max}** and prolonged T_{1/2} **compared to treprostinil**

Prolonged duration of pharmacodynamic activity compared to treprostinil

“Local” pharmacodynamic effect

- Achieves the same pharmacodynamic (vasodilatory) effect at significantly lower systemic concentrations, compared with IV treprostinil

Effect on hallmark histopathologic features in the sugen/hypoxia model

- Remodeling effect, superior to other prostanoids

Reduced incidence of cough (guinea pig model)

Highlights of Preclinical Data

Preclinical data demonstrate the following features of treprostinil palmitil:

Improved pharmacokinetics

- Prolonged lung residence time
- Lower systemic Cmax and prolonged T1/2 compared to treprostinil

Prolonged duration of pharmacodynamic activity compared to treprostinil

“Local” pharmacodynamic effect

- Achieves the same pharmacodynamic (vasodilatory) effect at significantly lower systemic concentrations, compared with IV treprostinil


Effect on hallmark histopathologic features in the sugen/hypoxia model

- Remodeling effect, superior to other prostanoids


Reduced incidence of cough (guinea pig model)

Improved Pharmacokinetics:

Prolonged Lung Residence, Lower Cmax, Prolonged Half Life (Rat and Dog)



F.G. Leifer et al. (2018) Drug Res 68: 1-10



M. R. Corboz, et al. (2017) J. Pharmacol. Exp. Ther. 363:348-357.

Highlights of Preclinical Data

Preclinical data demonstrate the following features of treprostinil palmitil:

Improved pharmacokinetics

- Prolonged lung residence time
- Lower systemic Cmax and prolonged T1/2 compared to treprostinil

Prolonged duration of pharmacodynamic activity compared to treprostinil

“Local” pharmacodynamic effect

- Achieves the same pharmacodynamic (vasodilatory) effect at significantly lower systemic concentrations, compared with IV treprostinil


Effect on hallmark histopathologic features in the sugen/hypoxia model

- Remodeling effect, superior to other prostanoids

Reduced incidence of cough (guinea pig model)


Prolonged Duration of Pharmacodynamic Activity

(Telemetered Rat, Hypoxic Challenge)



Prolonged Duration of Pharmacodynamic Activity

(Telemetered Dog, Hypoxic Challenge)



Average right ventricular pulse pressure (RVPP) response to hypoxia challenge in two dogs following inhalation of different doses of TP; the 'wash out' period between doses was ≥ 7 days. Shown at $t=0$ is the no-treatment baseline average (values were taken before dosing).

Highlights of Preclinical Data

Preclinical data demonstrate the following features of treprostinil palmitil:

Improved pharmacokinetics

- Prolonged lung residence time
- Lower systemic Cmax and prolonged T1/2 compared to treprostinil

Prolonged duration of pharmacodynamic activity compared to treprostinil

“Local” pharmacodynamic effect

- **Achieves the same pharmacodynamic (vasodilatory) effect at significantly lower systemic concentrations, compared with IV treprostinil**

Effect on hallmark histopathologic features in the sugen/hypoxia model


- Remodeling effect, superior to other prostanoids

Reduced incidence of cough (guinea pig model)

Local Pharmacodynamic Effect


Pulmonary Vasodilator Activity at Significantly Lower Plasma Treprostinil Concentrations (Rat)

Rats challenged with the thromboxane mimetic U46619



Local Pharmacodynamic Effect

Pulmonary Vasodilator Activity at Significantly Lower Plasma Treprostinil Concentrations (Dog)



Local Pharmacodynamic Effect

Continuously Inhaled vs. Continuously Infused (IV) Treprostinil in Sheep

“

...prostacyclin analogs are more potent pulmonary vasodilators when delivered by aerosol than when given intravenously. Systemic hemodynamic effects are also minimized by aerosol administration of the drug, making this approach especially appealing. Duration of action of prostacyclin is short, requiring an unrealistic frequency of administration for clinical use (6, 33), but development of analogs (e.g., treprostinil) or formulations that are long acting could make this approach feasible.”

J Appl Physiol 99: 2363–2368, 2005.

First published September 1, 2005; doi:10.1152/jappphysiol.00083.2005.

Potent effects of aerosol compared with intravenous treprostinil on the pulmonary circulation

Brett L. Sandifer,^{1,3} Kenneth L. Brigham,^{1,2,3} E. Clinton Lawrence,^{1,3} David Mottola,⁴ Chris Cuppels,^{1,2} and Richard E. Parker^{1,2}

¹Division of Pulmonary, Allergy, and Critical Care, ²Center for Translational Research in the Lung, and ³McKelvey Center for Lung Transplantation, Emory University School of Medicine, Atlanta, Georgia; and ⁴United Therapeutics Corporation, Research Triangle Park, North Carolina

Submitted 24 January 2005; accepted in final form 20 July 2005

“We speculate that this may result from treprostinil stimulated production of vasodilatory mediators from pulmonary epithelium.”

Highlights of Preclinical Data

Preclinical data demonstrate the following features of treprostinil palmitil:

Improved pharmacokinetics

- Prolonged lung residence time
- Lower systemic Cmax and prolonged T1/2 compared to treprostinil

Prolonged duration of pharmacodynamic activity compared to treprostinil

“Local” pharmacodynamic effect

- Achieves the same pharmacodynamic (vasodilatory) effect at significantly lower systemic concentrations, compared with IV treprostinil

Effect on hallmark histopathologic features in the sugen/hypoxia model

- **Remodeling effect, superior to other prostanoids**

Reduced incidence of cough (guinea pig model)

Treprostinil Palmitil Inhalation Solution


Improved Both Hemodynamics and Vascular Remodeling in the Rat Sugen-Hypoxia Model of PAH

* significantly different from the naïve group


significantly different from the vehicle group

¹Fulton Index is a commonly used quantitative measure of the degree of right ventricular hypertrophy
Am. J. Res. Crit. Care Med. 2020: 201, A2094. Poster was not presented due to COVID19.
Data on file, Insmid Inc.


Improves Pulmonary Arterial Pressure




Reduces Wall Thickening



Improves Fulton Index¹




Reduces Muscularization



TPIP Demonstrated Superior Effect

on PA Pressure and RV Hypertrophy in the Rat Sugen-Hypoxia Model of PAH




* significantly different from Normoxic control (Nx)

significantly different from Vehicle Group


¹Fulton Index is a commonly used quantitative measure of the degree of right ventricular hypertrophy: it is the weight ratio of right ventricle/(left ventricle + septum)

TPIP Demonstrated Superior Effect on Vascular Remodeling in the Sugen-Hypoxia Rat Model of PAH

Wall Thickness



Muscularization




Data on file, Insmad Inc.

* significantly different from Normoxic control group
significantly different from Vehicle group
* and # p < 0.05
** and ## p < 0.01
*** and ### p < 0.001

TPIP at High Dose

Shows Superior Effect Overall in the Sugden-Hypoxia Rat Model for PAH



Highlights of Preclinical Data

Preclinical data demonstrate the following features of treprostinil palmitil:

Improved pharmacokinetics

- Prolonged lung residence time
- Lower systemic Cmax and prolonged T1/2

Prolonged duration of pharmacodynamic activity

“Local” pharmacodynamic effect

- Achieves the same pharmacodynamic (vasodilatory) effect at significantly lower systemic concentrations, compared with IV treprostinil


Effect on hallmark histopathologic features in the sugen/hypoxia model

- Remodeling effect, superior to other prostanoids

Reduced incidence of cough (guinea pig model)

Reduced Incidence of Cough

Threshold to Induce Cough Increased 7-Fold (guinea pig model)



*Significantly different from vehicle.

**Treprostinil molar equivalent doses; TP doses are 2.3, 12.8 and 35.8 µg/kg.

Data on file, Inmed Inc.




Product Differentiation

Treprostinil Palmitil Inhalation Solution Phase 1 Study

Plasma PK in Healthy Volunteers

Treprostinil Plasma PK Summary		
Cohort 1 (n = 6)	TPIS 85 µg	Tyvaso 54 µg
C _{max} (ng/mL)	0.089	0.958
AUC ₀₋₂₄ (ng*h/mL)	0.614	0.872
T _{max} (h)	1.02	0.258
T _{1/2} (h)	5.69	0.485



Han et al. (2016) Single-Dose Pharmacokinetics of C16TR for Inhalation (INS1009) vs Treprostinil Inhalation Solution. ERS International Congress, September 3-7, London, United Kingdom, abs 3505 (poster PA2303).

Treprostinil Palmitil Inhalation Powder (TPIP)

a Potentially Differentiated Prostanoid

If the characteristics suggested by the preclinical studies are borne out in the clinical development program, TPIP could represent a differentiated product

Convenience: portable, dry powder inhaler

QD or BID dosing (prolonged lung residence time, prolonged half life, prolonged duration of pharmacodynamic activity)

Improved safety/tolerability

- Systemic tolerability (lower C_{max})
 - Local tolerability (reduced cough)
-


Greater vasodilatory pharmacodynamic effect at lower systemic exposures (sustained lung residence time takes advantage of the local effect seen in multiple animal models)

Potential disease modifier – may be able to benefit the histopathologic process, whereas Tyvaso, due to transient local exposure, cannot



Path Forward

Next Steps for TPIP Development



Phase 1 Underway

SAD/MAD Study

Objective

Establish PK and tolerability profile

Top line results expected in Q1 2021

Initiate Phase 2a in Early 2021

Exploratory, single dose study

Objective

Demonstrate proof of mechanism, understand PK/PD relationship, confirm tolerability

Results expected in 2021

Initiate Phase 2b in H2 2021

Objective

Demonstrate hemodynamic effect size, inform design for pivotal program

Kevin C. Mange, M.D., MSCE
SVP, Clinical Development

Colleen Silk
General Manager, Brensocatib




Brensocatib

Reversible, Oral DPP1 Inhibitor

Brensocatib

NCFBE to Serve as 'Launch Pad' for Neutrophil-mediated Portfolio



NCFBE
Program
WILLOW +
ASPEN



Serious Unmet Need Conditions



- Cystic Fibrosis
- Granulomatosis with Polyangiitis
- Alpha-1 Antitrypsin Deficiency

Larger Markets



- Inflammatory Bowel Disease*
- Rheumatoid Arthritis*
- Lupus Nephritis*
- COPD/Asthma†
- Oncology

Research Pipeline



- External collaborations
- Insmed expertise

NCFBE, non-CF bronchiectasis

** Preclinical data generated*

†AstraZeneca has exercised its option to develop brensocatib in COPD & asthma through Phase 2b. Further development subject to agreement on commercial terms with Insmed.



Brensocatib

Clinical Development

Bronchiectasis: Pathogenesis

Bronchiectasis Vicious Cycle*

INFLAMMATION

Persistent inflammatory response, causing lung damage

AIRWAY DESTRUCTION

An environmental insult or disease process leads to bronchial wall destruction and dilation



BACTERIAL COLONIZATION

Increased susceptibility to chronic bacterial infection and colonization

ABNORMAL MUCOCILIARY CLEARANCE

Structural damage impairs protective mucociliary clearance

* All references to bronchiectasis related to our WILLOW and ASPEN clinical studies, potential launch of brensocatib and global prevalence refer to non-cystic fibrosis bronchiectasis.

WILLOW Study Results Underscore Potential of Brensocatib



The NEW ENGLAND
JOURNAL of MEDICINE



ORIGINAL ARTICLE FREE PREVIEW

Phase 2 Trial of the DPP-1 Inhibitor Brensocatib in Bronchiectasis

James D. Chalmers, M.B., Ch.B., Ph.D., Charles S. Haworth, M.B., Ch.B., M.D., Mark L. Metersky, M.D., Michael R. Loebinger, B.M., B.Ch., Ph.D., et al., for the WILLOW Investigators*

September 7, 2020
DOI: 10.1056/NEJMoa2021713


Abstract

BACKGROUND Patients with bronchiectasis have frequent exacerbations that are thought to be related to neutrophilic inflammation. The activity and quantity of neutrophil serine proteases, including neutrophil elastase, are increased in the sputum of patients with bronchiectasis at baseline and increase further during exacerbations. Brensocatib (INS1007) is an oral reversible inhibitor of dipeptidyl peptidase 1 (DPP-1), an enzyme responsible for the activation of neutrophil serine proteases.

METHODS In a phase 2, randomized, double-blind, placebo-controlled trial, we randomly assigned, in a 1:1:1 ratio, patients with bronchiectasis who had had at least two exacerbations in the previous year to receive placebo, 10 mg of brensocatib, or 25 mg of brensocatib once daily for 24 weeks. The time to the first exacerbation (primary end point), the rate of exacerbations (secondary end point), sputum neutrophil elastase activity, and safety were assessed.

RESULTS Of 256 patients, 87 were assigned to receive placebo, 82 to receive 10 mg of brensocatib, and 87 to receive 25 mg of brensocatib. The 25th percentile of the time to the first exacerbation was 67 days in the placebo group, 134 days in the 10-mg brensocatib group, and 96 days in the 25-mg brensocatib group. Brensocatib treatment prolonged the time to the first exacerbation as compared with placebo ($P=0.03$ for 10-mg brensocatib vs. placebo; $P=0.04$ for

Brensocatib Phase 2 WILLOW Study Design



- **Primary objective:**

- Time to first exacerbation over 24 weeks in patients with non-cystic fibrosis bronchiectasis (NCFBE)

- **Secondary objectives:**





- Pulmonary exacerbation rate over 24 weeks
- Change in QOL-Bronchiectasis questionnaire respiratory symptoms domain over 24 weeks
- Change in post-bronchodilator ppFEV₁ over 24 weeks
- Change in sputum NE activity from pretreatment to on-treatment

25 mg brensocatib vs. placebo

Rate of Exacerbations by Patient Subgroups

10 mg brensocatib vs. placebo


Chalmers et al, ERS International
Congress 2020, 7-9 September. RCT4135
BSI, bronchiectasis severity index; FEV₁,
forced expiratory volume in 1 second
^aBulgaria/Poland




Dose Response Effect Seen Consistently Across Sputum Neutrophil Serine Proteases (NSPs)

Mean (SE) Change from Baseline to
Week 24 (log10 [mg/mL])


Neutrophil Elastase




Cathepsin G



Proteinase 3



WILLOW Safety Data



no. (%)	Placebo (n=85)	Brensocatib 10 mg (n=81)	Brensocatib 25 mg (n=89)
TEAE resulting in study discontinuation	3 (3.5)	3 (3.7)	4 (5.5)
TEAE resulting in treatment discontinuation	9 (10.6)	6 (7.4)	6 (6.7)
Serious TEAE	19 (22.4)	11 (13.6)	10 (11.2)
Serious TEAEs in ≥ 3% of patients in any group			
Infective exacerbation of bronchiectasis	9 (10.6)	5 (6.2)	4 (4.5)
Pneumonia	3 (3.5)	0	4 (4.5)
Any TEAE	67 (78.8)	75 (92.6)	74 (83.1)
TEAEs in ≥ 10% of patients in any group			
Cough	10 (11.8)	15 (18.5)	12 (13.5)
Headache	3 (3.5)	8 (9.9)	12 (13.5)
Sputum increased	6 (7.1)	9 (11.1)	9 (10.1)
Dyspnea	2 (2.4)	3 (3.7)	9 (10.1)
Infective exacerbation of bronchiectasis	9 (10.6)	5 (6.2)	4 (4.5)
Diarrhea	9 (10.6)	5 (6.2)	3 (3.4)



Brensocatib

Phase 3 Development Program

ASPEN Study: Single Trial Designed to Support Global Filing



Breakthrough Therapy Designation for
brensocatib in non-cystic fibrosis
bronchiectasis (NCFBE)

Robust efficacy endpoints, supported by
WILLOW results

Integrates feedback from Health Authorities

ASPEN Phase 3 Study Design

Primary Endpoint:


- Rate of pulmonary exacerbations over the 52-week treatment period

Key Secondary Endpoints:

- Time to first pulmonary exacerbation over the 52-week treatment period
- Proportion of subjects who remain exacerbation-free over the 52-week treatment period
- Change in FEV1

Powered at 90% to detect 30% reduction in exacerbation rate

- Type I error: 0.01 (two-sided)
- P-value = less than 0.01



ASPEN Timeline Considerations



36 Countries



~480 sites



6 Fast Track Countries



Targeting WILLOW study sites



Brensocatib

Commercial Outlook

Brensocatib Strategy Builds on Successful First Launch in NCFBE



ASPEN Phase 3 trial for non-cystic fibrosis bronchiectasis (NCFBE)

Prepare for successful global launch in NCFBE

Advance clinical programs for cystic fibrosis

Evaluate other neutrophil-modulated diseases for future research & label expansion

An Approved NCFBE Treatment to Reduce Exacerbations

Could Address Unmet Needs Across Stakeholders



Patients

- Symptoms of cough, dyspnea, fatigue interfere with daily life
- Fear of exacerbations negatively impacts emotional and social well-being
- Frustrated with time-consuming therapies that are not specific to BE



Healthcare Providers

- Lack of efficacious anti-inflammatory option
- Treatments that improve patient QOL and prevent exacerbations seen as largest unmet need
- Recognize current treatments lack BE-specific clinical trial data



Payers

- Treatment option with BE indication, greater efficacy, and reduction in exacerbations seen as unmet needs

“


It's like waiting for a hurricane – there's a constant level of fear and anxiety of the unknown. When will an exacerbation hit?

”

- BRONCHIECTASIS PATIENT


Bronchiectasis is a Globally Prevalent Disease

Preliminary Data Suggest Significant Global Need



**Weycker, et al. Prevalence and incidence of NCFBE among US adults in 2013. Chronic Respiratory Disease. 2017; **Estimates suggest broadly similar per capita prevalence in EU5 as in US; ◇Asia-Pacific rates 3X to ~10X higher than those in the US; Zhou, YM et al. The prevalence and risk factors of bronchiectasis in residents aged 40 years old and above in seven cities in China. 2013 † EU5 comprised of France, Germany, Italy, Spain and the United Kingdom*

Current Prevalence Estimates May Underestimate Global Patient Population, Given Underdiagnosis of NCFBE



1. Agusti, A., Calverley, P.M., Celli, B. et al. Characterisation of COPD heterogeneity in the ECLIPSE cohort. *Respir Res.* 2010; 11: 122
2. Ni, Y., Shi, G., Yu, Y. et al. Clinical characteristics of patients with chronic obstructive pulmonary disease with co-morbid bronchiectasis: a systematic review and meta-analysis. *Int J Chron Obstruct Pulmon Dis.* 2015; 10: 1465–1475
3. Kang HR, Choi GS, Park SJ, et al. The effects of bronchiectasis on asthma exacerbation. *Tuberc Respir Dis (Seoul).* 2014;77(5):209-214.

Well Positioned to Support a Successful Global Launch in NCFBE

Uniquely positioned to commercialize brensocatib, if approved, given the synergies that exist between NCFBE and NTM lung disease



Well-developed US launch capabilities with ARIKAYCE

Experience with patient finding and activation

Geographic and target HCP overlap with current US Sales structure

EU/Japan teams mobilized for launch success


Advocacy and patient support




Brensocatib

Additional Opportunities

Strategy to Build a Neutrophil-mediated Portfolio



NCFBE
Program



Serious Unmet Need Conditions



- Cystic Fibrosis
- Granulomatosis with Polyangiitis
- Alpha-1 Antitrypsin Deficiency

Larger Markets



- Inflammatory Bowel Disease*
- Rheumatoid Arthritis*
- Lupus Nephritis*
- COPD/Asthma†
- Oncology

Research Pipeline



- External collaborations
- Insmed expertise

* Preclinical data generated

†AstraZeneca has exercised its option to develop brensocatic in COPD & asthma through Phase 2b. Further development subject to agreement on commercial terms with Insmed.

Cystic Fibrosis and GPA Identified for Further Clinical Development

	Cystic Fibrosis	GPA
MOA Applicability	Clear role for neutrophils and NE in CF; Chronic bacterial infections of CF are neutrophil-mediated, and very elevated neutrophil counts in sputum are seen in CF patients	Two distinct MOAs could be applicable: (1) Inhibits activity of NSPs, (2) Reduces expression of membrane-bound PR3
Value Proposition	Potential to fill a need for CF-specific anti-inflammatory therapy, a category not currently met by marketed products An important additive therapy (on top of CFTR modulators) for CF patients who experience any pulmonary exacerbations	Potential for additional treatment option for GPA patients during induction for a more efficacious response, or as an alternative option for sustained remission
Providing a Roadmap for Development from CF to GPA	Previously approved products (CFTR modulators) and support from the CF Foundation Therapeutics Development Network	Previously approved product (Rituxan) and phase 3 development of avacopan

Ongoing Investigator-Sponsored Clinical Study for Treatment of Severe COVID-19 Infections

Could brensocatib be used to treat COVID-19?

Currently under investigation in multi-site UK investigator-initiated research (IIR) study

Double-blind, prospective, placebo-controlled, randomized trial

Up to 300 patients; 10 sites

Led by University of Dundee's Professor James Chalmers, MBChB, Ph.D.

Prioritized by UK's National Institute for Health Research

Potential Data read-out: **H1 2021**



Kevin C. Mange, M.D., MSCE
SVP, Clinical Development

Neil Hughes
General Manager, Head of EMEA

Roger Adsett
Chief Operating Officer



Kevin C. Mange, M.D., MSCE
SVP, Clinical Development



ARIKAYCE

Front Line Program

Not for promotional use

ARIKAYCE® (amikacin liposomal inhalation suspension)

Indication and Use

- LIMITED POPULATION: ARIKAYCE® is indicated in adults, who have limited or no alternative treatment options, for the treatment of *Mycobacterium avium* complex (MAC) lung disease as part of a combination antibacterial drug regimen in patients who do not achieve negative sputum cultures after a minimum of 6 consecutive months of a multidrug background regimen therapy. As only limited clinical safety and effectiveness data for ARIKAYCE are currently available, reserve ARIKAYCE for use in adults who have limited or no alternative treatment options. This drug is indicated for use in a limited and specific population of patients.
- This indication is approved under accelerated approval based on achieving sputum culture conversion (defined as 3 consecutive negative monthly sputum cultures) by Month 6. Clinical benefit has not yet been established. Continued approval for this indication may be contingent upon verification and description of clinical benefit in confirmatory trials.
- **Limitation of Use:** ARIKAYCE has only been studied in patients with refractory MAC lung disease defined as patients who did not achieve negative sputum cultures after a minimum of 6 consecutive months of a multidrug background regimen therapy. The use of ARIKAYCE is not recommended for patients with non-refractory MAC lung disease.

WARNING: RISK OF INCREASED RESPIRATORY ADVERSE REACTIONS

ARIKAYCE has been associated with an increased risk of respiratory adverse reactions, including hypersensitivity pneumonitis, hemoptysis, bronchospasm, and exacerbation of underlying pulmonary disease that have led to hospitalizations in some cases.

First-in-Disease Therapy for MAC Lung Disease



MAC lung disease is a rare, progressive, and chronic condition that can cause severe, permanent damage to the lungs



Symptoms often worsen over time, including chronic cough, dyspnea, fatigue, fever, weight loss, and chest pain



Caused by bacteria in the environment and is more likely to affect those with a history of lung conditions, like bronchiectasis or COPD

ARIKAYCE

Novel, inhaled, once-daily formulation of liposomal amikacin taken with a nebulizer



Pulmovance™ liposomal technology delivers drug directly to the lungs

Prolongs release of amikacin in the lungs while limiting systemic exposure

Complementary portfolio

Meaningful overlap in patient populations, with over 60% of bronchiectasis patients also having an NTM infection

ARIKAYCE Front Line Indication – Trial Summary

ARISE (INS-415)

Psychometric validation study

Subjects: 100


Start: Q4 2020

ENCORE (INS-416)

Double-blind PMR trial


Subjects: 250

Start: Q4 2020



Front Line Indication

Registration Program



ARISE

Primary Objective

Demonstrate reliability, validity and responsiveness of the PRO/symptom scores

Secondary Objective

Demonstrate effect of ARIKAYCE on culture conversion, time to culture conversion

ENCORE

Primary Endpoint

Change from Baseline to Month 13 (one month off treatment) in respiratory symptom score

Key Secondary Endpoint

Proportion of subjects achieving durable culture conversion at Month 15 (3 months off treatment)

*Azithromycin (AZI), Ethambutol (ETH)

**Patients will be enrolled into separate open label extension study

Developing a Symptom Score for Front Line Indication

Leveraging Existing Instruments to Expedite Work

Respiratory Symptom Score

Quality of Life- Bronchiectasis (QoL-B)
Patient Reported Outcome (PRO)
Questionnaire

Fatigue Symptom Score

PROMIS-Fatigue Short Form 7a
Questionnaire

In general, FDA requires description (validation) of questionnaires (PROs) to be used to support efficacy in the context of use (population of interest)

Completed Concept Elicitation work with 25 patients with newly diagnosed MAC disease

Supports specific questions in respiratory domain in QoL-B, and in PROMIS-Fatigue are meaningful / understandable to these patients and are likely the basis of computing symptom scores pending completion of Cognitive Interview work later this year.

Developing a Symptom Score for Front Line Indication

Lessons Learned from Prior Drug Development Programs

Describing longitudinal properties of questionnaires / PROs is required by FDA, in general

Global Blood Therapeutics incorporated describing longitudinal properties into Phase 3 study (HOPE study) for their new PRO in sickle cell disease

ARISE & ENCORE will evaluate symptom scores and other endpoints after completion of three-drug treatment regimen vs two-drug treatment regimen

ARISE will evaluate longitudinal properties as a separate study that will complete while ENCORE is enrolling

Provides for ability to adjust what/how items are used to compute final version of symptom scores in ENCORE while preserving integrity of ENCORE

Neil Hughes

General Manager, Head of EMEA



ARIKAYCE a Blueprint for Future Success

2019 **\$136M** net sales

Q2 2020 **\$42.5M**

Engaged HCP & patient population

Positive reimbursement trends

- Engaged healthcare providers
- Educated patient population
- Focus on payers to ensure reimbursement

ARIKAYCE was one of the Top 10 most successful non-oncology rare disease launches

We plan to repeat history with our growing pipeline



ARIKAYCE Added to International Treatment Guidelines for Treatment of MAC Lung Disease



We help the world breathe®
PULMONARY • CRITICAL CARE • SLEEP




Guidelines are a globally recognized standard for the prevention, diagnosis and treatment of NTM lung disease

Addition of ARIKAYCE to new standard treatment regimen for adult patients with limited or no treatment options who failed to convert to a negative sputum culture after at least six months of treatment

Treatment should be continued for 12 months after culture conversion


Solid EU Infrastructure in Place to Support Launch*



*If approved

Commercial Approach

Meaningful Opportunity in Europe



**Source: Internal analysis of published NTM epidemiology, primary market research with treating HCPs, and anonymized patient level claims data in US*

[†] EU5 comprised of France, Germany, Italy, Spain and the United Kingdom

ARIKAYCE Received Positive CHMP Opinion

July 24, 2020

Committee for Medicinal Products for Human Use (CHMP) of the European Medicines Agency has adopted a positive opinion recommending ARIKAYCE for the treatment of NTM lung infections caused by MAC in non-CF Patients with limited treatment options



Prevalence of NTM in EU May be Greater Than Previously Reported

DISPATCHES

Prevalence of Nontuberculous Mycobacterial Pulmonary Disease, Germany, 2009–2014

Felix C. Ringshausen, Dirk Wagner,
Andrés de Roux, Roland Diel, David Hohmann,
Lennart Hickstein, Tobias Welte,
Jessica Rademacher

We analyzed routine statutory health insurance claim data to determine prevalence of nontuberculous mycobacterial pulmonary disease in Germany. Documented prevalence rates of this nonnotifiable disease increased from 2.3 to 3.3 cases/100,000 population from 2009 to 2014. Prevalence showed a strong association with advanced age and chronic obstructive pulmonary disease.

health insurance (Table 1) (7). Because this study was based on anonymous routine data, institutional review board approval and patient consent were not required. We obtained anonymous health claims data from the Health Risk Institute health services research database, which contains a subset of ≈7 million persons covered by statutory health insurance. This database has been validated and shows good overall accuracy in documenting the German population in terms of illness, death, and drug usage; in addition, the number of insured persons in the database remains high over time (8). We established data characteristics by adjustments for age and sex according to the



EUROPEAN RESPIRATORY *journal*

FLAGSHIP SCIENTIFIC JOURNAL OF ERS

Early View

Original article


Identification of potentially undiagnosed patients with nontuberculous mycobacterial lung disease using machine learning applied to primary care data in the UK

Orla M. Doyle, Roald van der Laan, Marko Obradovic, Peter McMahon, Flora Daniels, Ashley Pitcher, Michael R. Loebinger

Please cite this article as: Doyle OM, van der Laan R, Obradovic M, *et al.* Identification of potentially undiagnosed patients with nontuberculous mycobacterial lung disease using machine learning applied to primary care data in the UK. *Eur Respir J* 2020; in press (<https://doi.org/10.1183/13993003.00045-2020>).

...while anticipated EU indication slightly wider than only refractory MAC

EU Launch Strategy



Insmed Projected EMEA Country Footprint

Specific Field Force Structures Reflect Country-Specific Patient Pathways and NTM Management Protocols

Country Key

- Regional HQ – No Local Commercial Org
- Insmed Commercial Organization
- Under Review - Insmed vs Distributor




Customer facing FTEs grow as country reimbursement comes online*

FTEs	2020	2021	2022
Field Based Medical	14	18	21
Field Based Commercial	10	21	29

*Projections

Anticipated EU Launch Sequence*






Anticipated timelines shown

Insmed has been accepted to participate in an accelerated reimbursement process being piloted by the Dutch Health Care Institute (ZIN). While ARIKAYCE is currently well advanced in the accelerated review process, there is no guarantee ARIKAYCE will ultimately qualify for accelerated reimbursement. In the event that ARIKAYCE does complete this process, we anticipate to launch in the Netherlands according to the traditional timeline shown above. Not for promotional use

*If approved

Germany

First Anticipated Launch & Largest European Market



Four "3-up" territories


MSL, RDM & PSP nurse cover identical geography to allow for smooth collaboration between the different parts of the Field Force

Analytics Enable Focused Commercialization Model in Germany

Geographic analysis based on:

- Centers of Excellence
- Areas of high NTM prevalence
- Existing clinical experience with ARIKAYCE

Split into priority focus areas and
“white space”



Roger Adsett

Chief Operating Officer




ARIKAYCE

Japan Launch Update

Not for promotional use


Commercial Approach: Expansion to Japan



**Source: Internal analysis of published NTM epidemiology, primary market research with treating HCPs, and anonymized patient level claims data in US*


† EU5 comprised of France, Germany, Italy, Spain and the United Kingdom

On Track for Anticipated Approval, Mid-Year 2021 Japan Launch



Anticipated timelines shown

“Cost Plus” Approach for Japanese Pricing



Nuances of Japanese Market



Prescriptions are limited to **two-week supply** for the first year drug is on the market



Co-pay market: approved drugs reimbursed at fixed rates



Physician offices would exclusively train patients on use of ARIKAYCE with Lamira device

Building a Strong Presence in Japan

28 People with Japanese Affiliate



Engaging Major Medical Associations



The Research Institute of Tuberculosis,
Japan Anti-tuberculosis Association



The Japanese Association for
Infectious Diseases



一般社団法人
日本結核・非結核性抗酸菌症学会
THE JAPANESE SOCIETY FOR TUBERCULOSIS AND NONTUBERCULOUS MYCOBACTERIOSIS


Sales Territories in Japan*



Sales Director
East Japan



Sales Director
West Japan




**If approved*

A close-up photograph of a microscope with blue nitrile gloves. The microscope is focused on a sample on a slide, with a bright light illuminating the area. The background is blurred.

Sara Bonstein
Chief Financial Officer

Financial Update & Closing Remarks

Pipeline



* As a condition of accelerated approval, Insmed plans to conduct an additional clinical study to support full approval.

Financial Strength to Achieve Our Objectives

\$641.9M

as of 6/30/20

Cash and Cash
Equivalents

>3 Years

as of 9/30/20

Estimated Cash Runway

101M*

as of 6/30/20

Common Shares
Outstanding

**Excludes stock options, unvested RSUs and shares underlying
outstanding convertible notes*




Transformational Growth Strategy

World Class Team

Research &
Development Expertise

Global Commercial Presence

Business Development Strength





Questions & Answers



**We are committed to bringing forth
technologies and medicines in
therapeutic areas with the greatest
potential to make a difference in
patients' lives**

Thank You

Investor Relations
investor.relations@insmed.com
646-351-0954